

Birds of the Tambopata Reserve (Explorer's Inn Reserve)

Theodore A. Parker, III, Paul K. Donahue and Thomas S. Schulenberg

	Habitats	Foraging	Sociality	Abundance	Evidence
TINAMIDAE (9)					
<i>Tinamus major</i>	Ft	T	S	F	t
<i>T. guttatus</i>	Fh,Ft	T	S	F	t
<i>Crypturellus cinereus</i>	Ft	T	S	C	t
<i>C. soui</i>	Ft	T	S	F	t
<i>C. undulatus</i>	Z,Ft	T	S	C	t
<i>C. variegatus</i>	Fh	T	S	R	t
<i>C. bartletti</i>	Ft	T	S	U	t
<i>C. parvirostris*</i>	C	T	S	R	si
<i>C. atrocapillus</i>	Fh?	T	S	R	?
PODICIPEDIDAE (1)					
<i>Podiceps dominicus</i>	L	W	S	R	si
PHALACROCORACIDAE (1)					
<i>Phalacrocorax brasiliensis</i>	L,R	W	S,G	U	ph
ANHINGIDAE (1)					
<i>Anhinga anhinga</i>	L,R	W	S	U	ph
ARDEIDAE (15)					
<i>Pilherodias pileatus</i>	S,Lm	W	S	F	ph,t
<i>Ardea cocoi</i>	S,Rm,Lm	W	S	U	ph
<i>Casmerodius albus</i>	S,Lm	W	S	R	ph
<i>Bubulcus ibis</i>	S,C,Lm	T	S,G	U	ph
<i>Egretta caerulea*</i>	S,Rm	W	S	R	si
<i>E. thula</i>	S,Rm,Lm	W	S	U	ph
<i>Butorides striatus</i>	Lm,M	W	S	U	ph,t
<i>Agamia agami</i>	Lm,Fsm,Fs	W	S	R	ph
<i>Nycticorax nycticorax*</i>	Lm	W	S	R	si
<i>Cochlearius cochlearius</i>	Lm	W	S	U	ph
<i>Tigrisoma lineatum</i>	Lm,Fsm,Fs	W	S	U	ph,t
<i>Zebrilus undulatus</i>	Fs,Fsm	T,W	S	R	ph
<i>Ixobrychus involucris*</i>	M	T,W	S	R(Ms)	si
<i>I. exilis</i>	M	T,W	S	R(M?)	t
<i>Botaurus pinnatus</i>	M	T,W	S	R(Ms)	ph

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
CICONIIDAE (2)					
<i>Mycteria americana</i>	S	W	G,S	U	si
<i>Jabiru mycteria</i>	S	W	S	R	si
THRESKIORNITHIDAE (2)					
<i>Mesembrinibis cayennensis</i>	Fs,Lm	T,W	S	U	ph
<i>Ajaia ajaja</i>	S	W	S	R	si
ANHIMIDAE (2)					
<i>Anhima cornuta</i>	M,S	T	S	F	ph,t
<i>Chauna torquata</i>	S	T	S	R(Ms)	ph
ANATIDAE (3)					
<i>Dendrocygna</i> sp.	O	W	G	R	si
<i>Neochen jubata</i>	S	T	S	R	si
<i>Cairina moschata</i>	R,L,Fs	W	S	U	ph
CATHARTIDAE (4)					
<i>Sarcoramphus papa</i>	Ft	T	S	U	si
<i>Coragyps atratus</i>	S,Rm,Ft	T	S,G	U	si
<i>Cathartes aura</i>	S,Rm,Ft	T	S	R	si
<i>Cathartes melambrotus</i>	Z,Ft,Fh	T	S	F	si
ACCIPITRIDAE (27)					
<i>Chondrohierax uncinatus</i> *	Lm	Sc,U	S	R	si
<i>Gampsonyx swainsonii</i>	Rm,Fe	C,A	S	R	si
<i>Elanoides forficatus</i>	Ft	A	S,G	R	si
<i>Leptodon cayanensis</i>	Ft,Fh	C,Sc	S	U	t
<i>Harpagus bidentatus</i>	Fh	Sc	S	U	t
<i>Ictinia plumbea</i>	Z,Ft	A	S,G	U	si
<i>Rostrhamus sociabilis</i>	M	T	S	R	si
<i>R. hamatus</i>	Lm,M	U,T	S	R	si
<i>Accipiter poliogaster</i>	Ft	Sc,C	S	R	si
<i>A. bicolor</i>	Ft	U,Sc	S	R	si
<i>A. superciliosus</i>	Ft,Fh	C,Sc	S	R	t
<i>A. striatus</i>	O	C,Sc	S	R(Ms)	si
<i>Buteo albonotatus</i>	Ft	C,T	S	R	si
<i>B. platypterus</i> *	Ft,Fe	C,Sc	S	R(Mn)	si
<i>B. magnirostris</i>	Z,Rm,Fe	T,C	S	F	t
<i>B. brachyurus</i>	Ft	A,C	S	R	si

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>Asturina nitida*</i>	Z,Fe	T,C	S	R	si
<i>Leucopternis kuhli</i>	Ft,Fh	C	S	R	t
<i>L. schistacea</i>	Fs,Fsm,Ft	T,Sc	S	U	t
<i>Buteogallus urubitinga</i>	Rm,Lm,S	T,U	S	U	ph,t
<i>Busarellus nigricollis</i>	Fs,Fsm,Lm	T,U	S	R	t
<i>Morphnus guianensis</i>	Ft,Fh	Sc,C	S	R	si
<i>Harpia harpya</i>	Ft,Fh	Sc,C	S	R	ph
<i>Spizastur melanoleucus</i>	Ft	C,A?	S	R	si
<i>Spizaetus ornatus</i>	Ft,Fh	C,T	S	R	t
<i>Spizaetus tyrannus</i>	Z,Ft	Sc,C	S	U	t
<i>Geranoospiza caerulescens</i>	Ft,Fs	T,C	S	U	t
PANDIONIDAE (1)					
<i>Pandion haliaetus</i>	L,R	W	S	U(Mn)	si
FALCONIDAE (9)					
<i>Herpetotheres cachinnans</i>	Ft,Fh,	C,T	S	U	t
<i>Micrastur mirandollei</i>	Ft,Fh	Sc,U	S	R	t
<i>M. ruficollis</i>	Ft,Fh	Sc,U	S	U	t
<i>M. gilvicollis</i>	Ft,Fh	Sc,U	S	U	t
<i>Daptrius ater</i>	Rm,S,Z	T,C	S,G	U	t
<i>D. americanus</i>	Ft,Fh	C	G	F	t
<i>Milvago chimachima</i>	Rm	T,C	S	R	si
<i>Falco deiroleucus*</i>	Rm,Fe	C,A	S	R	si
<i>Falco rufigularis</i>	Rm,C	A	S	F	t
CRACIDAE (4)					
<i>Ortalis guttata</i>	Z,Ft	C	G	U	ph,t
<i>Penelope jacquacu</i>	Ft,Fh	C,T	S,G	F	ph,t
<i>Aburria pipile</i>	Ft,Z	C	S	U	ph,t
<i>Mitu tuberosa</i>	Ft,Fh,Fsm	T	S	R	ph,t
PHASIANIDAE (1)					
<i>Odontophorus stellatus</i>	Ft,Fh	T	G	U	t
OPISTHOCOMIDAE (1)					
<i>Opisthocomus hoazin</i>	Lm	U,C	G	C	ph,t
ARAMIDAE (1)					
<i>Aramus guarauna</i>	Lm,M	W	S	R	t

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
PSOPHIIDAE (1)					
<i>Psophia leucoptera</i>	Fh,Ft	T	G	U	t
RALLIDAE (8)					
<i>Rallus nigricans</i>	M	T,W	S	R	si
<i>Aramides cajanea</i>	Lm,Fsm,Fs	T	S	F	t
<i>Anurolimnas castaneiceps</i>	C	T	S	R	t
<i>Lateralus exilis</i>	M	T,W	S	U	t
<i>L. melanophaius</i>	M	T,W	S	U	t
<i>Neocrex erythrops*</i>	M	T	S	R	si
<i>Porphyryla martinica</i>	M	T,W	S	U	si
<i>P. flavirostris</i>	M	T,W	S	U(M?)	sp
HELIORNITHIDAE (1)					
<i>Heliornis fulica</i>	L	W	S	U	ph
EURYPYGIDAE (1)					
<i>Eurypyga helias</i>	Fsm,Fs,S	T,W	S	U	t
JACANIDAE (1)					
<i>Jacana jacana</i>	M	T,W	S,G	U	t
CHARADRIIDAE (4)					
<i>Hoploxypterus cayanus</i>	S	T	S	F	si
<i>Vanellus sp.*</i>	O,C,S	T	S	R	si
<i>Charadrius collaris</i>	S	T	S	U	t
<i>Pluvialis dominica</i>	S	T	S,G	R(Mn)	si
SCOLOPACIDAE (9)					
<i>Tringa solitaria</i>	S	T	S	U(Mn)	si
<i>T. flavipes</i>	S	T	S	R(Mn)	si
<i>T. melanoleuca</i>	S	T	S	R(Mn)	si
<i>Actitis macularia</i>	S	T	S	F(Mn)	t
<i>Calidris fuscicollis*</i>	S	T	S	R(Mn)	si
<i>C. melanotos</i>	S	T	S,G	R(Mn)	si
<i>C. himantopus*</i>	S	T	S,G	R(Mn)	si
<i>Tryngites subruficollis</i>	S	T	G	R(Mn)	si
<i>Bartramia longicauda</i>	C,S	T	S	R(Mn)	si
LARIDAE (4)					
<i>Larus pipixcan*</i>	L	W	S	R(Mn)	si
<i>Phaetusa simplex</i>	R,L	W,A	S	U	t

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>Sterna superciliaris</i>	R,L	W,A	S	F	t
<i>Sterna nilotica</i> *	R	W,A	S	R(Mn)	si
RHYNCHOPIDAE (1)					
<i>Rynchops niger</i>	R,L	W	S,G	R	ph,t
COLUMBIDAE (9)					
<i>Columba speciosa</i> *	O(Rm)	C	S	R	si
<i>C. cayennensis</i>	Lm,Z	U,C	S,G	C	ph,t
<i>C. subvinacea</i>	Ft	C	S	U	t
<i>C. plumbea</i>	Ft,Fh	C	S	C	t
<i>Columbina talpacoti</i>	C,Rm,Lm	T	S,G	U	si
<i>C. picui</i>	C,Rm	T	S,G	U(Ms?)	si
<i>Claravis pretiosa</i>	Z	T	S	R	t
<i>Leptotila rufaxilla</i>	Z,C	T	S	F	ph,t
<i>Geotrygon montana</i>	Ft,Fh	T	S	F	ph,t
PSITTACIDAE (19)					
<i>Ara ararauna</i>	A,Fs,Ft	C	G	U	ph,t
<i>A. macao</i>	Ft,Fh	C	S,G	F	ph,t
<i>A. chloroptera</i>	Ft,Fh	C	S,G	F	ph,t
<i>A. severa</i>	Z,Ft	C	S,G	F	ph,t
<i>A. manilata</i>	Ft	C	G	R	ph,t
<i>A. couloni</i>	Ft	C	S,G	R	t
<i>Aratinga leucophthalmus</i>	Z,Ft	C	G	U	ph,t
<i>A. weddellii</i>	Z,Ft	C	G	C	ph,t
<i>Pyrrhura rupicola</i>	Ft,Fh	C	G	F	t
<i>Forpus sclateri</i>	Ft,Fh	C	S	U	t
<i>Brotogeris cyanoptera</i>	Z,Ft,Fh	C	G	C	ph,t
<i>B. sanctithomae</i> *	Z	C	G	R	si
<i>Touit huetii</i>	?(O)	C	G	R	t
<i>Nannopsittaca dachilleae</i>	Ft	C	S	R	t
<i>Pionites leucogaster</i>	Ft,Fh	C	G	F	t
<i>Pionopsitta barrabandi</i>	Fh,Ft	Sc,C	S,G	F	t
<i>Pionus menstruus</i>	Ft	C	S,G	C	t
<i>Amazona ochrocephala</i>	Ft,Z	C	S	U	t
<i>A. farinosa</i>	Ft,Fh	C	S,G	C	t

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
CUCULIDAE (13)					
<i>Coccyzus cinereus</i> *	Ft,W,Z	C	M,S	R(Ms)	si
<i>C. erythrophthalmus</i> *	Fh	C	M,S	R(Mn)	si
<i>C. americanus</i>	Ft,Fh	C,Sc	M,S	U(Mn)	si
<i>C. melacoryphus</i>	Z,Ft	C	S,M	U(Ms)	si
<i>Piaya cayana</i>	Z,Ft,Fh	Sc,C	S,M	F	t
<i>P. melanogaster</i>	Ft,Fh	C	S,M	U	t
<i>P. minuta</i>	Lm	U,Sc	S	U	t
<i>Crotophaga major</i>	Rm,Lm,Fs	U,Sc	G	F(W)	t
<i>C. ani</i>	C,Lm	U,T	G	F	t
<i>Tapera naevia</i>	C	T,U	S	R	t
<i>Dromococcyx phasianellus</i>	Ft	T,U	S	R	t
<i>D. pavoninus</i>	Ft,B	T,U	S	R	t
<i>Neomorphus geoffroyi</i>	Fh	T	S	R	si
STRIGIDAE (8)					
<i>Otus choliba</i>	Z	Sc,C	S	R	t
<i>O. watsonii</i>	Ft,Fh	U,C	S	C	t
<i>Lophotrix cristata</i>	Ft,Fh	C	S	U	t
<i>Pulsatrix perspicillata</i>	Ft	C,T	S	U	t
<i>Glaucidium hardyi</i>	Ft,Fh	C	S	F	t
<i>G. brasilianum</i>	Z	C,Sc	S	U	t
<i>Ciccaba virgata</i>	Ft,Fh	C	S	U	sp
<i>C. huhula</i>	Ft,Fh?	C	S	R	?
NYCTIBIIDAE (4)					
<i>Nyctibius grandis</i>	Ft	C,A	S	U	t
<i>N. aethereus</i>	Ft	C,A	S	R	t
<i>N. griseus</i>	Z,Ft,Fe	C,A	S	U	t
<i>N. bracteatus</i> *	Ft	?	S	R	si
CAPRIMULGIDAE (8)					
<i>Lurocalis semitorquatus</i>	O(Ft)	A	S	R	t
<i>Chordeiles rupestris</i>	S	A	G	C	si
<i>C. minor</i>	Fe	A	G	R(Mn)	si
<i>Nyctidromus albicollis</i>	Z,C	A	S	F	t
<i>Nyctiphrynus ocellatus</i>	Ft,B	A,Sc	S	U	t
<i>Hydropsalis climacocerca</i>	Rm,S	A	S	F	t

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>Caprimulgus parvulus</i>	Ft	A,Sc	S	R(Ms)	t
<i>C. sericocaudatus</i>	Ft,Fh	A,C	S	R(Ms?)	t
APODIDAE (8)					
<i>Streptoprocne zonaris</i>	O(Ft,Fh)	A	G	F	si
<i>Cypseloides</i> sp.	O(Ft)	A	G	R(M?)	si
<i>Chaetura chapmani</i>	O(Ft)	A	G,M	R(M?)	si
<i>Chaetura cinereiventris</i>	O(Ft)	A	G,M	C(M?)	t
<i>C. egregia</i>	O(Ft)	A	G,M	U(Ms?)	t?
<i>C. brachyura</i>	O(Fe)	A	G,M	U	t
<i>Panyptila cayennensis</i>	O(Ft)	A	S,M	U	t
<i>Reinarda squamata</i>	O(A,L,R)	A	S,G	C	t
TROCHILIDAE (18)					
<i>Glaucis hirsuta</i>	Z,Ft	U,Sc	S	U	t
<i>Threnetes leucurus</i>	Z,Ft	U	S	R	sp,t
<i>Phaethornis philippi</i>	Fh	U	S	F	t
<i>P. hispidus</i>	Z,Ft	U	S	F	sp,t
<i>P. ruber</i>	Ft,Fh,B	U	S	C	t
<i>Campylopterus largipennis</i>	Ft,Z	U,Sc	S	R	si
<i>Florisuga mellivora</i>	Ft,Fh	C	S	U	si
<i>Anthracothorax nigricollis</i>	Rm,Lm	C	S	U	si
<i>Lophornis chalybea</i>	Ft,W	C	S	R	si
<i>Popelairia</i> sp.	C	C	S	R	si
<i>Thalurania furcata</i>	Ft,Fh	U,C	S	F	t
<i>Hylocharis cyanus</i>	Ft	C	S	F	t
<i>Chrysuronia oenone</i>	C,W	C	S	R	si
<i>Amazilia lactea</i>	Z,Ft	C	S	R	si
<i>Polyplancta aurescens</i>	Ft	Sc,C	S	U	t
<i>Heliostyris aurita</i>	Ft,Fh	Sc,C	S	U	si
<i>Heliomaster longirostris</i>	Z,Rm	C	S	R	si
<i>Calliphlox amethystina</i>	W	C	S	R	si
TROGONIDAE (6)					
<i>Pharomachrus pavoninus</i>	Fh,Ft	C	S	U	t
<i>Trogon melanurus</i>	Ft,Fh	C	S	C	t
<i>T. viridis</i>	Ft,Fh	C,Sc	S,M	C	t
<i>T. collaris</i>	Ft,Fh	Sc,U	S,M	F	t

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>T. curucui</i>	Ft,Z	C	S,M	U	t
<i>T. violaceus</i>	Ft,Fh	C	S,M	U	t
ALCEDINIDAE (5)					
<i>Ceryle torquata</i>	Rm,Lm	W	S	F	t
<i>Chloroceryle amazona</i>	Rm,Lm	W	S	F	t
<i>C. americana</i>	Fsm,Rm,Lm	W	S	U	si
<i>C. inda</i>	Fsm,Lm	W	S	U	si
<i>C. aenea</i>	Fsm,Lm	W	S	U	si
MOMOTIDAE (3)					
<i>Electron platyrhynchum</i>	Ft,Fh	C,Sc	S	C	t
<i>Baryphengus martii</i>	Ft	Sc,C	S	F	t
<i>Momotus momota</i>	Ft,Fh	Sc,C	S	U	t
GALBULIDAE (4)					
<i>Brachygalba albogularis</i>	Z	C,A	S	R	ph,t
<i>Galbula cyanescens</i>	Ft,Fo	U,Sc	S,M	F	sp,t
<i>G. dea</i>	Fh	C	S,M	U	t
<i>Jacamerops aurea</i>	Ft,Fh	C,Sc	S	U	t
BUCCONIDAE (10)					
<i>Notharchus macrorhynchos</i>	Ft,Fh	C	S	U	t
<i>N. ordii</i>	Ft	C	S	R	ph
<i>Bucco macrodactylus</i>	Z	Sc,C	S	R	si
<i>Nystalus striolatus</i>	Ft,Fh	C,	S	U	t
<i>Malacoptila semicincta</i>	Ft,Fh	U	S	U	t
<i>Nonnula ruficapilla</i>	Ft,Z,B	U,Sc	S	U	sp,t
<i>Monasa nigrifrons</i>	Ft,Z,	C,Sc	G,M	C	t
<i>M. morphoeus</i>	Ft,Fh	C,Sc	G,M	C	t
<i>M. flavirostris</i>	Ft	C	S,M	R	t
<i>Chelidoptera tenebrosa</i>	Rm,Z,Ft	C,A	S	F	t
CAPITONIDAE (3)					
<i>Capito niger</i>	Ft,Fh	C	S,M	C	t
<i>Eubucco richardsoni</i>	Ft	C	S,M	F	t
<i>E. tucinkae</i>	Z,Ft	C	S,M	R	t
RAMPHASTIDAE (8)					
<i>Aulacorhynchus prasinus</i>	Ft	C	S	U	t
<i>Pteroglossus castanotis</i>	Ft,Z	C	G	F	t

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>P. inscriptus</i>	Ft	C	G	R	t
<i>P. mariae</i>	Ft,Fh	C	G	F	t
<i>Pteroglossus beauharnaesii</i>	Ft,Fh	C	G	U	t
<i>Selenidera reinwardtii</i>	Ft,Fh	C,Sc	S	F	t
<i>Ramphastos culminatus</i>	Ft,Fh	C	S,G	C	t
<i>R. cuvieri</i>	Ft,Fh	C	S,G	C	t
PICIDAE (16)					
<i>Picumnus rufiventris</i> *	Z	Sc	S,M	R	si
<i>P. aurifrons</i>	Ft,Fh	C	S,M	U	sp
<i>Chrysoptilus punctigula</i>	Z	Sc,C	S	R	t
<i>Piculus leucolaemus</i>	Ft	C	M	U	t
<i>P. chrysochloros</i>	Ft,Fh	C,Sc	S,M	U	t
<i>Celeus elegans</i>	Ft,Fs	C,Sc	S,M	R	t
<i>C. grammicus</i>	Fh,Ft	C	S,M	F	t
<i>C. flavus</i>	Ft	C,Sc	S,G	U	t
<i>C. spectabilis</i>	Z,Ft,B	U,Sc	S,M	U	ph,t
<i>C. torquatus</i>	Ft,Fh	C,Sc	S	R	t
<i>Dryocopus lineatus</i>	Z,Ft	Sc,C	S	F	t
<i>Melanerpes cruentatus</i>	Ft,Fh,Fo	C	S,M	C	t
<i>Veniliornis passerinus</i>	Z	Sc,C	S,M	F	t
<i>V. affinis</i>	Ft,Fh	C	M	F	t
<i>Campephilus melanoleucus</i>	Ft,Z	C,Sc	S	F	t
<i>C. rubricollis</i>	Ft,Fh	U,Sc	S	F	t
DENDROCOLAPTIDAE (16)					
<i>Dendrocincla fuliginosa</i>	Ft	U,Sc	S,M,A	F	t
<i>D. merula</i>	Ft,Fh	U,Sc	S,A	U	t
<i>Deconychura longicauda</i>	Ft,Fh	Sc	S,M	U	t
<i>Sittasomus griseicapillus</i>	Ft	U,C	M	F	t
<i>Glyphorhynchus spirurus</i>	Ft,Fh	U,Sc	S,M	C	sp
<i>Nasica longirostris</i>	Lm,Fs,Ft	Sc,C	S	U	t
<i>Dendrexetastes rufigula</i>	Ft,Fh	C,Sc	S,M	F	t
<i>Xiphocolaptes promeropirhynchus</i>	Fh	U,C	S,M	R	t
<i>Dendrocolaptes certhia</i>	Ft,Fh	Sc,C	S,M,A	F	t
<i>D. picumnus</i>	Ft,Fh	U,Sc	S,A	U	t
<i>Xiphorhynchus picus</i>	Z	Sc	S	R	t

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>X. obsoletus</i>	A?	U,Sc	S,M	R	t
<i>X. spixii</i>	Ft,Fh	U,Sc	M	C	sp,t
<i>X. guttatus</i>	Ft,Fh	Sc,C	S,M	C	t
<i>Lepidocolaptes albolineatus</i>	Ft,Fh	C	M	F	t
<i>Campylorhamphus trochilirostris</i>	Ft,B	U	S,M	U	sp
FURNARIIDAE (28)					
<i>Furnarius leucopus</i>	Z,Rm	T	S	F	t
<i>Synallaxis cabanisi</i>	B	U	S	R	t
<i>S. albescens</i>	Rm,C	U	S	R(Ms?)	sp
<i>S. albigularis</i>	Rm,C	U	S	F	t
<i>S. gujanensis</i>	Z	U	S	F	t
<i>S. rutilans</i>	Ft,Fh	T,U	S	F	t
<i>Cranioleuca gutturata</i>	Ft	C,Sc	M	U	t
<i>Thripophaga fusciceps</i>	Ft	U,Sc	S,M	R	sp
<i>Berlepschia rikeri</i>	A	C	S	R	t
<i>Hyloctistes subulatus</i>	Ft	Sc,C	S,M	U	sp,t
<i>Ancistrops strigilatus</i>	Ft,Fh	Sc,C	M	C	sp,t
<i>Simoxenops ucayale</i>	Ft,B,Z	U	M,S	U	sp
<i>Philydor erythrocerus</i>	Fh,Ft	Sc	M	F	t
<i>P. pyrrhodes</i>	Ft,Fh	Sc,U	S,M	U	t
<i>P. rufus</i>	Z	C	S,M	U	t
<i>P. erythropterus</i>	Ft,Fh	C	M	F	t
<i>P. ruficaudatus</i>	Ft	Sc,C	M	R	t
<i>Automolus infuscatus</i>	Ft,Fh	U	M	C	sp,t
<i>A. dorsalis</i>	Ft,Z,B	Sc	S,M	R	t
<i>A. rubiginosus</i>	Ft,B	U	S	R	t
<i>A. ochrolaemus</i>	Ft,Fh	U	S,M	F	ph,t
<i>A. rufipileatus</i>	Z,Ft	U	S,M	F	t
<i>A. melanopezus</i>	Ft,B	U,Sc	S,M	U	sp,t
<i>Xenops milleri</i>	Ft,Fh	C	M	F	t
<i>X. rutilans</i>	Ft,Fh	C	M	R	t
<i>X. minutus</i>	Ft,Fh	U,Sc	M	F	sp,t
<i>Sclerurus mexicanus*</i>	Ft	T	S	R	si
<i>S. caudacutus</i>	Ft,Fh	T	S	F	t

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
FORMICARIIDAE (52)					
<i>Cymbilaimus lineatus</i>	Ft,Fh	Sc,C	S,M	F	t
<i>C. sanctaemariae</i>	Ft,B	Sc,C	S,M	F	ph,t
<i>Frederickena unduligera</i>	Fh	U	S	R	t
<i>Taraba major</i>	Z,Lm	U	S	U	t
<i>Thamnophilus doliatus</i>	Rm,C	U	S,M	U	t
<i>T. aethiops</i>	Ft,Fh	U	S	F	t
<i>T. schistaceus</i>	Ft,Fh	Sc	M	C	t
<i>T. amazonicus</i>	Fs,Lm	Sc	S	R	t
<i>Pygiptila stellaris</i>	Ft,Fh	C,Sc	M	C	t
<i>Thamnomanes ardesiacus</i>	Ft,Fh	U	M	F	t
<i>T. schistogynus</i>	Ft	U,Sc	M	C	sp,t
<i>Myrmotherula brachyura</i>	Ft,Fh	C	M	C	t
<i>M. sclateri</i>	Ft,Fh	C	M	F	t
<i>M. surinamensis</i>	Lm	U,Sc	S,M	F	t
<i>M. longicauda</i>	Z	U	S	R	t?
<i>M. hauxwelli</i>	Ft,Fh	U	S,M	F	sp,t
<i>M. leucophthalma</i>	Ft,Fh	U	M	F	sp,t
<i>M. ornata</i>	Ft,B	U,Sc	M	F	sp,t
<i>M. axillaris</i>	Ft,Fh	U,Sc	M	C	sp,t
<i>M. iheringi</i>	Ft,B	U	M	U	sp,t
<i>M. longipennis</i>	Fh	U,Sc	M	F	t
<i>M. menetriesii</i>	Ft,Fh	Sc,C	M	C	t
<i>Dichrozona cincta</i>	Fh	T	S	R	t
<i>Microrhophias quixensis</i>	Ft,B	Sc	M	R	t
<i>Drymophila devillei</i>	Ft,B	Sc	S,M	F	t
<i>Terenura humeralis</i>	Ft	C	M	U	t
<i>Cercomacra cinerascens</i>	Ft,Fh	C,Sc	S,M	C	t
<i>C. nigrescens</i>	Z,B	U	S	U	ph,t
<i>C. serva</i>	Fh,Ft,Fo	U	S	R	ph,t
<i>C. manu</i>	Z,B	U	S	R	t
<i>Myrmoborus leucophrys</i>	Ft	U	S	F	sp,t
<i>M. myotherinus</i>	Ft,Fh	U	S,A	C	t
<i>Hypocnemis cantator</i>	Ft,B	U	S,M	C	sp,t
<i>Hypocnemoides maculicauda</i>	Fs,Lm,Fsm	U,T	S	F	sp

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>Percnostola lophotes</i>	Z,B	T,U	S	F	sp,t
<i>Sclateria naevia</i>	Fs,Fsm,Lm	T	S	U	t
<i>Myrmeciza hemimelaena</i>	Ft,Fh	U,T	S	C	t
<i>M. hyperythra</i>	Fs,Lm	T,U	S	F	t
<i>M. goeldii</i>	Ft,B	U	S,A	U	sp,t
<i>M. atrothorax</i>	Z	U	S	U	t
<i>Gymnopithys salvini</i>	Ft,Fh	U	A	F	sp
<i>Rhegmatorhina melanosticta*</i>	Fh	U	A	R	si
<i>H. punctulata</i>	Ft	U	S	R	ph
<i>H. poecilinota</i>	Ft,Fh	U	S,A	U	ph,t
<i>Phlegopsis nigromaculata</i>	Ft	T,U	S,A	U	ph,t
<i>Chamaeza nobilis</i>	Ft	T	S	R	t
<i>Formicarius colma</i>	Fh	T	S	F	ph,t
<i>F. analis</i>	Ft	T	S,A	C	ph,t
<i>F. rufifrons</i>	Ft	T	S	R	t
<i>Hylopezus berlepschi</i>	Z	T	S	R	t
<i>Myrmothera campanisona</i>	Ft,Fh	T	S	F	t
<i>Conopophaga peruviana</i>	Ft	U,T	S	R	t
COTINGIDAE (7)					
<i>Iodopleura isabellae</i>	Ft	C	S	U	t
<i>Porphyrolaema porphyrolaema*</i>	Ft	C	M	R	si
<i>Lipaugus vociferans</i>	Ft,Fh	Sc,C	S	C	t
<i>Cotinga maynana</i>	Ft,Z	C	S	F	t
<i>C. cayana</i>	Fh	C	S	U	si
<i>Gymnoderus foetidus</i>	Ft	C	S	F	si
<i>Querula purpurata</i>	Fh,Ft	C	G	C	t
PIPRIDAE (10)					
<i>Schiffornis major</i>	Fs,Lm	U	S	U	sp
<i>S. turdinus</i>	Fh	U	S	U	t
<i>Piprites chloris</i>	Ft,Fh	C	M	F	t
<i>Xenopipo atronitens</i>	Z	U	S	R	sp
<i>Tyrannetes stolzmanni</i>	Ft,Fh	Sc	S	C	t
<i>Machaeropterus pyrocephalus</i>	Ft,B	Sc,C	S	F	sp,t
<i>Pipra coronata</i>	Fh	U,Sc	S	U	t
<i>P. fasciicauda</i>	Ft	U,Sc	S	F	sp,t

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>P. rubrocapilla</i>	Fh	U,Sc	S	U	sp,t
<i>P. chloromeros</i>	Ft,Fh	U,Sc	S	F	t
TYRANNIDAE (90)					
<i>Zimmerius gracilipes</i>	Ft,Fh	C	S,M	F	t
<i>Ornithion inerme</i>	Ft,Fh	C	S,M	F	sp,t
<i>Camptostoma obsoletum</i>	Z	C	S,M	U	t
<i>Sublegatus modestus</i>	W,C	Sc,C	S,M	R(M?)	si
<i>S. obscurior</i>	Z,C	Sc,C	S,M	R	t
<i>Phaeomyias murina</i>	C	C	S,M	U	t
<i>Tyrannulus elatus</i>	Ft,Fh	C,Sc	S,M	F	t
<i>Myiopagis gaimardii</i>	Ft,Fh	C	M	C	sp,t
<i>M. caniceps</i>	Ft	C	M	R	t
<i>M. viridicata</i>	Ft,Fh,Z	Sc,C	M	U(Ms)	t
<i>Elaenia spectabilis</i>	Z	C,Sc	S,M	U(Ms)	sp
<i>E. parvirostris</i>	Z	C,Sc	S,M	R(Ms)	si
<i>E. strepera*</i>	B,Ft	U,Sc	S?	R(Ms)	si
<i>Inezia inornata</i>	Z	C	G,M	F(Ms)	sp,t
<i>Pseudocolopteryx acutipennis</i>	W,M	U	S	R(Ms?)	sp
<i>Euscarthmus meloryphus</i>	Z	U	S	R(Ms)	t
<i>Mionectes olivaceus</i>	Ft,Fh	C,Sc	M	R	si
<i>M. oleagineus</i>	Ft,Fh	U,C	M	F	sp,t
<i>M. macconnelli</i>	Ft,Fh	U,Sc	M	R	t?
<i>Leptopogon amaurocephalus</i>	Ft	U,Sc	M,S	F	t
<i>Corythopis torquata</i>	Ft,Fh	T	S	F	t
<i>Myiornis ecaudatus</i>	Ft	C,Sc	S	F	t
<i>Poecilatriccus albifacies</i>	B,Ft	U,Sc	M,S	U	sp
<i>Hemitriccus flammulatus</i>	Ft,B	Sc	S	F	sp,t
<i>H. zosterops</i>	Ft,Fh	Sc	S	F	sp,t
<i>H. iohannis</i>	Z	Sc,C	S	U	t
<i>H. striaticollis</i>	Z	U	S	R(Ms)	sp
<i>Todirostrum latirostre</i>	Z	U	S	F	ph,t
<i>T. maculatum</i>	Z	C,Sc	S,M	F	t
<i>T. chrysocrotaphum</i>	Ft,Fh	C	S,M	F	t
<i>Ramphotrigon megacephala</i>	Z,Ft,B	Sc,U	S,M	F	sp,t
<i>R. fuscicauda</i>	Ft,B	Sc	S	U	ph,t

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>R. ruficauda</i>	Ft,Fh	Sc	S,M	F	t
<i>Rhynchocyclus olivaceus</i>	Ft	U,Sc	M	U	ph,t
<i>Tolmomyias assimilis</i>	Ft,Fh	C	M	F	t
<i>T. poliocephalus</i>	Ft,Z	C	S,M	F	t
<i>T. flaviventris</i>	Z	C,Sc	S,M	F	t
<i>Platyrinchus coronatus</i>	Ft	Sc,U	S	F	sp
<i>P. platyrhynchos</i>	Fh	U,Sc	S	R	t
<i>Onychorhynchus coronatus</i>	Ft,Fh,Fsm	U,Sc	S,M	U	t
<i>Terentotriccus erythrurus</i>	Ft,Fh	U,Sc	S,M	U	t
<i>Myiophobus fasciatus</i>	C,Z	U	S,M	U	t
<i>Contopus borealis</i>	Rm	C	S	R(Mn)	si
<i>C. virens</i>	C,Z	C,A	S	U(Mn)	si
<i>C. cinereus</i>	Z	C,A	S	R(M?)	si
<i>Empidonax alnorum</i>	C,Z	U	S	R(Mn)	sp
<i>Lathrotriccus eulerei</i>	Ft,Z,B	U,Sc	S	F	t
<i>Cnemotriccus fuscatus</i>	Z	U,Sc	S	U	t
<i>Pyrocephalus rubinus</i>	Rm,C	C,A	S	F(Ms)	si
<i>Ochthoeca littoralis</i>	Rm,S	T,A	S	F	t
<i>Muscisaxicola fluviatilis</i>	Rm,S	T	S	U	si
<i>Knipolegus sp.*</i>	Lm,M	U	S	R(Ms?)	si
<i>Hymenops perspicillata*</i>	M	U	S	R(Ms)	si
<i>Fluvicola pica</i>	Lm,M	U,T	S	R(Ms?)	si
<i>Colonia colonus</i>	Rm,Z	C	S	R(Mm?)	t
<i>Satrapa icterophrys</i>	Rm	Sc	S,M	R(Ms)	sp
<i>Attila cinnamomeus</i>	Ft	Sc	S	R	t
<i>A. bolivianus</i>	Ft	Sc,C	S	U	t
<i>A. spadiceus</i>	Ft,Fh	Sc,C	S,M	F	t
<i>Casiornis rufa</i>	Ft	C	M	R(Ms)	sp
<i>Rhytipterna simplex</i>	Ft,Fh	Sc,C	S,M	C	t
<i>Laniocera hypopyrra</i>	Ft,Fh	Sc,C	S,M	F	t
<i>Sirystes sibilator</i>	Ft,Fh	C,Sc	S,M	F	t
<i>Myiarchus tuberculifer</i>	Ft	C	S	R(N?)	t
<i>M. swainsoni</i>	Z,Ft	C	M	U(Ms)	t?
<i>M. ferox</i>	Z,C	Sc,C	S,M	F	sp,t
<i>M. tyrannulus</i>	Ft,Fh	C	S,M	F(Ms)	t

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>Pitangus lictor</i>	M	U	S	R	t
<i>P. sulphuratus</i>	Rm,C	U,Sc	S	F	t
<i>Megarhynchus pitangua</i>	C,Z	Sc,C	S	F	t
<i>Myiozetetes cayanensis</i>	Lm,M	U	S	F	t
<i>M. similis</i>	C,Rm	U,Sc,C	S,G	C	t
<i>M. granadensis</i>	Z,C	Sc,C	S,G	F	t
<i>M. luteiventris</i>	Ft	C	S,G	U	t
<i>Myiodynastes maculatus</i>	Ft	Sc,C	M	U(Ms)	si
<i>M. luteiventris</i>	Ft,Z,W	C	S,M	U(Mn)	si
<i>Legatus leucophaeus</i>	Ft,Fo,Lm	C	S	F	ph,t
<i>Empidonomus varius</i>	Ft	C	M	R(Ms)	si
<i>E. aurantioatrocristatus</i>	Ft,Fh,Fo	C	S,M	F(Ms)	si
<i>Tyrannopsis sulphurea</i>	Lm,A	C	S	U	t
<i>Tyrannus melancholicus</i>	Rm,C	C,A	S	F	t
<i>T. savanna</i>	Rm,Lm	C	G	U(Ms?)	si
<i>T. tyrannus</i>	Ft,Rm,Lm	C	G	C(Mn)	si
<i>Pachyramphus castaneus</i>	Z	C	M	R	t?
<i>P. polychopterus</i>	Z,C	C	S,M	F	sp,t
<i>P. marginatus</i>	Ft,Fh	C	M	F	t
<i>P. minor</i>	Ft,Fh	C	M	U	t
<i>P. validus</i>	Fh	C	M	R	si
<i>Tityra cayana</i>	Ft	C	S	F	si
<i>T. semifasciata</i>	Ft,Fh	C	S	U	t
<i>T. inquisitor</i>	Ft	C	S	U	si
HIRUNDINIDAE (13)					
<i>Tachycineta albiventer</i>	R	A	S,G	C	t
<i>T. leucorhoa*</i>	C,O	A	G	R(Ms)	si
<i>Phaeoprogne tapera</i>	R	A	G,S	U	t
<i>Progne chalybea</i>	R	A	S,G	R	si
<i>P. modesta*</i>	R,L	A	G	R(Ms)	si
<i>Notiochelidon cyanoleuca</i>	R	A	G	R(Ms?)	si
<i>Atticora fasciata</i>	R,C	A	G	C	t
<i>Neochelidon tibialis*</i>	Fh	A	U	R	si
<i>Alopochelidon fucata*</i>	R,L	A	G	R(Ms)	si
<i>Stelgidopteryx ruficollis</i>	R,C	A	G	C	t

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>Riparia riparia</i>	L	A	G	R(Mn)	si
<i>Hirundo rustica</i>	R,C,L	A	G	F(Mn)	si
<i>Petrochelidon pyrrhonota</i>	L	A	G	R(Mn)	si
CORVIDAE (1)					
<i>Cyanocorax violaceus</i>	Z,Ft	Sc,C	G,M	C	t
TROGLODYTIDAE (7)					
<i>Campylorhynchus turdinus</i>	C,Z	Sc,C	S,M	F	t
<i>Thryothorus genibarbis</i>	Z,Ft	U	S	C	sp,t
<i>T. leucotis</i>	C	U	S	U	sp,t
<i>Troglodytes aedon</i>	C	U	S	U	t
<i>Microcerculus marginatus</i>	Ft,Fh	T	S	F	t
<i>Cyphorhinus arada</i>	Ft, Fh	T	S,M	F	t
<i>Donacobius atricapillus</i>	M,C	U	S	F	t
TURDIDAE (8)					
<i>Catharus fuscescens*</i>	Ft	T,U	S	R(Mn)	si
<i>C. minimus*</i>	Ft	T,U	S	R(Mn)	si
<i>C. ustulatus*</i>	Ft,Z	T,U	S	R(Mn)	si
<i>Turdus amaurochalinus</i>	Z,C,Ft	T,Sc,C	S	F(Ms?)	sp,t
<i>T. ignobilis</i>	Z,W,C	T,Sc,C	S,M	U	si
<i>T. lawrencii</i>	Ft	T,Sc,C	S	R	t
<i>T. hauxwelli</i>	Ft	T,Sc	S	U	t
<i>T. albicollis</i>	Ft,Fh	T,Sc	S	F	sp,t
VIREONIDAE (5)					
<i>Cyclarhis gujanensis</i>	Z	C	S,M	R	t
<i>Vireo olivaceus</i>	Ft,Fh,Z	C	M	C(Ms)	t
<i>Hylophilus thoracicus</i>	Ft	C	M	U	t
<i>H. hypoxanthus</i>	Ft,Fh	C	M	C	t
<i>H. ochraceiceps</i>	Fh	U	M	R	t
EMBERIZINAE (13)					
<i>Myospiza aurifrons</i>	S,C	T,U	S	C	t
<i>Volatinia jacarina</i>	C,M	T,U	S,G	R	si
<i>Sporophila schistacea</i>	Ft,B	U,Sc	S	R	sp
<i>S. lineola</i>	M,W,C	U	G,M	U(M?)	si
<i>S. luctuosa*</i>	W	U	G,M	R(Mm?)	si
<i>S. nigricollis</i>	W,C	U	G,M	R(Mm?)	si

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>S. caeruleascens</i>	M,C	T,U	G,M	C(Ms)	sp
<i>S. leucoptera</i> *	W,M	U	M	R(Ms)	si
<i>S. castaneiventris</i>	C	U,C	S,M	U	si
<i>Oryzoborus maximiliani</i>	M	U	S	R	t
<i>O. angolensis</i>	C,M	U	S	R	si
<i>Arremon taciturnus</i>	Ft,Fh	T,U	S	F	t
<i>Paroaria gularis</i>	Rm	U,Sc,C	S,M	F	sp
CARDINALINAE (7)					
<i>Pheucticus aureoventris</i> *	C	Sc	S	R(Ms?)	si
<i>P. ludovicianus</i> *	Ft	C	M	R(Mn)	si
<i>Caryothraustes humeralis</i>	Ft,Fh	C	M	R	si
<i>Pitylus grossus</i>	Ft	Sc	S,M	C	ph,t
<i>Saltator maximus</i>	Z,Ft,Fh	Sc,C	S,M	F	t
<i>S. coerulescens</i>	Z,C	U,Sc,C	S	F	t
<i>Cyanocopsa cyanoides</i>	Ft	U	S	F	t
THRAUPINAE (40)					
<i>Conothraupis speculigera</i>	Z	U,Sc	M	R(M?)	si
<i>Lamprospiza melanoleuca</i>	Fh,Ft	C	M	F	si
<i>Cissopis leveriana</i>	Z,C	Sc,C	S,M	F	t
<i>Thlypopsis sordida</i>	W,C,B	U	M	R(Ms?)	si
<i>Hemithraupis guira</i>	Ft	C	M	R	si
<i>H. flavicollis</i>	Ft,Fh	C	M	F	t
<i>Nemosia pileata</i> *	W	C	M	R(M?)	si
<i>Lanio versicolor</i>	Ft,Fh	Sc,C	M	F	t
<i>Tachyphonus cristatus</i>	Fh	C	M	U	si
<i>T. luctuosus</i>	Z,Ft,Fh	Sc,C	M	C	t
<i>Habia rubica</i>	Fh,Ft	U	G,M	C	t
<i>Piranga rubra</i> *	Ft	C	M	R(Mn)	si
<i>P. olivacea</i>	Ft	C	M	R(Mn)	si
<i>Ramphocelus nigrogularis</i>	Ft	Sc,C	G,M	C	t
<i>R. carbo</i>	C,Z,Ft	U,Sc,C	G,M	C	t
<i>Thraupis episcopus</i>	Z,C	Sc,C	S,M	U	si
<i>T. palmarum</i>	Ft,C	C	S,M	F	t
<i>Pipraeidea melanonota</i> *	C,W	C	M	R(Mm?)	si
<i>Euphonia laniirostris</i>	Z,C	Sc,C	S,M	U	t

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>E. musica</i> *	Fh	C	S	R(M?)	si
<i>E. chrysopasta</i>	Ft	Sc,C	S,M	F	t
<i>E. minuta</i>	Ft	C	M	U	t?
<i>E. xanthogaster</i>	Ft,Fh	U,Sc,C	M	F	t
<i>E. rufiventris</i>	Ft, Fh	Sc,C	S,M	C	t
<i>Chlorophonia cyanea</i>	Ft,Fh	C	M	R(M?)	t
<i>Tangara mexicana</i>	Ft	C	M,G	F	t
<i>T. chilensis</i>	Ft,Fh	Sc,C	G,M	C	t
<i>T. schrankii</i>	Ft,Fh	U,Sc,C	M	C	t
<i>T. xanthogastra</i>	Ft,Fh	C	M	R	si
<i>T. gyrola</i> *	Ft	C	G	R	si
<i>T. nigrocineta</i>	Ft,Fh	C	M	F	t
<i>T. velia</i>	Fh	C	M	U	t
<i>T. callophrys</i>	Fh	C	M	U	t
<i>Dacnis lineata</i>	Ft,Fh	C	M	C	t
<i>D. flaviventer</i>	Ft	C	S,M	U	t
<i>D. cayana</i>	Ft,Fh	C	M	C	si
<i>Chlorophanes spiza</i>	Ft,Fh	C	S,M	U	si
<i>Cyanerpes caeruleus</i>	Ft,Fh	C	S,G,M	C	si
<i>C. cyaneus</i> *	Ft	C	M	R	si
<i>Tersina viridis</i>	Z,Rm	C	G	U	t
PARULIDAE (6)					
<i>Dendroica striata</i> *	C,Z	C	M	R(Mn)	si
<i>Geothlypis aequinoctialis</i>	M,C	U	S	F	sp,t
<i>Oporornis agilis</i>	C	U	S	R(Mn)	sp
<i>Wilsonia canadensis</i> *	Ft	U	M	R(Mn)	si
<i>Basileuterus rivularis</i>	Ft,Fh,Fsm	T,U	S	F	t
<i>Coereba flaveola</i> *	C	Sc	M	R	si
ICTERIDAE (14)					
<i>Molothrus bonariensis</i>	Rm	T	G	R	si
<i>Scaphidura oryzivora</i>	Ft,Rm,S	T,C	S,G	F	t
<i>Clypicerus oseryi</i>	Ft	C	S,G	R	t
<i>Psarocolius decumanus</i>	Ft,Fh	C	G,M	F	t
<i>P. angustifrons</i>	Ft,Z	Sc,C	G,M	C	t
<i>Gymnostinops yuracares</i>	Ft,Fs	C	G,M	F	t

Habitats	
Fh	Upland Forest
Ft	Floodplain forest
Fs	Swamp forest
Fsm	Forest stream margins
Fo	Forest openings
Fe	Forest edges
Z	"Zabolo"
B	Bamboo
W	"Willow Bar"
C	Clearing
R	River
Rm	River margins
S	Shores
L	Lake
Lm	Lake margins
M	Marshes
A	Aguajales
O	Overhead
Foraging Position	
T	Terrestrial
U	Undergrowth
Sc	Subcanopy
C	Canopy
W	Water
A	Aerial
Sociality	
S	Solitary or in pairs
G	Gregarious
M	Mixed-species flocks
A	Army ant followers
Abundance	
C	Common
F	Fairly common
U	Uncommon
R	Rare
(M)	Migrant
(Mn)	Migrant from north
(Ms)	Migrant from south
(Mm)	Migrant from Andes
(W)	Wet season only
Evidence	
sp	Specimen
t	Tape
si	Species ID by sight
ph	Photo
*	recorded - 3 times

APPENDIX 3

	Habitats	Foraging	Sociality	Abundance	Evidence
<i>Cacicus cela</i>	Ft,Z	Sc,C	G,M	C	t
<i>C. haemorrhous</i>	Ft,Fh	Sc,C	S,G,M	U	t
<i>C. solitarius</i>	Z	U,Sc	S	F	t
<i>Agelaius xanthophthalmus</i>	M	U	S	R	t
<i>Icterus cayanensis</i>	Ft,C	C	S,M	F	t
<i>I. icterus</i>	Lm,M,C	U,Sc	S	U	ph,t
<i>Leistes superciliaris</i> *	C	T,U	S,G	R(Ms)	si
<i>Dolichonyx oryzivorus</i>	C	U	S,G	R(Mn)	si
Total:	572 species				
Resident Forest Species:	ca. 322 species				

* recorded 3 or fewer times, and only by sight (further verification required)